

K-theory: An Elementary Introduction

Mark Tomforde

October 14, 2020

Preliminaries

A **Hilbert space** is a vector space H with an inner product $\langle \cdot, \cdot \rangle$ that is complete with respect to the norm $\|x\| := \sqrt{\langle x, x \rangle}$.

$$B(H) := \{T : H \rightarrow H : T \text{ is linear and continuous}\}.$$

Preliminaries

A **Hilbert space** is a vector space H with an inner product $\langle \cdot, \cdot \rangle$ that is complete with respect to the norm $\|x\| := \sqrt{\langle x, x \rangle}$.

$$B(H) := \{T : H \rightarrow H : T \text{ is linear and continuous}\}.$$

For any $T \in B(H)$ there is a unique $T^* \in B(H)$, called the **adjoint**, for which

$$\langle T(x), y \rangle = \langle x, T^*(y) \rangle \text{ for all } x, y \in H.$$

$B(H)$ is a $*$ -algebra.

Preliminaries

A **Hilbert space** is a vector space H with an inner product $\langle \cdot, \cdot \rangle$ that is complete with respect to the norm $\|x\| := \sqrt{\langle x, x \rangle}$.

$$B(H) := \{T : H \rightarrow H : T \text{ is linear and continuous}\}.$$

For any $T \in B(H)$ there is a unique $T^* \in B(H)$, called the **adjoint**, for which

$$\langle T(x), y \rangle = \langle x, T^*(y) \rangle \text{ for all } x, y \in H.$$

$B(H)$ is a $*$ -algebra. Also:

$$K(H) := \{T \in B(H) : \overline{T(\text{Ball } H)} \text{ compact}\} = \overline{\{T \in B(H) : \text{rank}(T) < \infty\}}.$$

Fact $K(H) \triangleleft B(H)$. The **Calkin algebra** is $\mathcal{C}(H) := B(H)/K(H)$.

Preliminaries

A **Hilbert space** is a vector space H with an inner product $\langle \cdot, \cdot \rangle$ that is complete with respect to the norm $\|x\| := \sqrt{\langle x, x \rangle}$.

$$B(H) := \{T : H \rightarrow H : T \text{ is linear and continuous}\}.$$

For any $T \in B(H)$ there is a unique $T^* \in B(H)$, called the **adjoint**, for which

$$\langle T(x), y \rangle = \langle x, T^*(y) \rangle \text{ for all } x, y \in H.$$

$B(H)$ is a $*$ -algebra. Also:

$$K(H) := \{T \in B(H) : \overline{T(\text{Ball } H)} \text{ compact}\} = \overline{\{T \in B(H) : \text{rank}(T) < \infty\}}.$$

Fact $K(H) \triangleleft B(H)$. The **Calkin algebra** is $\mathcal{C}(H) := B(H)/K(H)$.

A **C^* -algebra** is a closed $*$ -subalgebra of $B(H)$.

Preliminaries

C^* -algebras have a connection with topology . . .

Recall:

$$C(X) := \{f : X \rightarrow \mathbb{C} : f \text{ is continuous}\}.$$

Preliminaries

C^* -algebras have a connection with topology . . .

Recall:

$$C(X) := \{f : X \rightarrow \mathbb{C} : f \text{ is continuous}\}.$$

Theorem: If A is a unital commutative C^* -algebra, then $A \cong C(X)$ for some compact Hausdorff space X .

Preliminaries

C^* -algebras have a connection with topology . . .

Recall:

$$C(X) := \{f : X \rightarrow \mathbb{C} : f \text{ is continuous}\}.$$

Theorem: If A is a unital commutative C^* -algebra, then $A \cong C(X)$ for some compact Hausdorff space X .

All topological information of X is encoded as algebraic information in $C(X)$, So “abelian C^* -algebras” are the same as “compact Hausdorff topological spaces”.

The study of C^* -algebras allows one to develop “noncommutative topology”.

Question What is K -theory (for Operator Algebras)?

Short Answer: A Homology Theory for C^* -algebras.

Question What is K -theory (for Operator Algebras)?

Short Answer: A Homology Theory for C^* -algebras.

Question Why do I, as an operator algebraist, care about K -theory?

Short Answer: It provides some of the most important invariants for C^* -algebras. These invariants allow you to show that particular C^* -algebras are different, ascertain knowledge about the C^* -algebra, and sometimes (perhaps surprisingly often) show two C^* -algebras are the same.

Question What is K -theory (for Operator Algebras)?

Short Answer: A Homology Theory for C^* -algebras.

Question Why do I, as an operator algebraist, care about K -theory?

Short Answer: It provides some of the most important invariants for C^* -algebras. These invariants allow you to show that particular C^* -algebras are different, ascertain knowledge about the C^* -algebra, and sometimes (perhaps surprisingly often) show two C^* -algebras are the same.

Question: What does the K stand for?

Answer: Grothendieck used the letter K to stand for “Klasse”, which means “class” in German (Grothendieck 's mother tongue).

Question What is K -theory (for Operator Algebras)?

Short Answer: A Homology Theory for C^* -algebras.

Question Why do I, as an operator algebraist, care about K -theory?

Short Answer: It provides some of the most important invariants for C^* -algebras. These invariants allow you to show that particular C^* -algebras are different, ascertain knowledge about the C^* -algebra, and sometimes (perhaps surprisingly often) show two C^* -algebras are the same.

Question: What does the K stand for?

Answer: Grothendieck used the letter K to stand for “Klasse”, which means “class” in German (Grothendieck’s mother tongue).

Question Where does K -theory (for Operator Algebras) come from?

Short Answer: Algebraic/Differential Topology.

Topological K -theory \subseteq Operator K -theory \subseteq Algebraic K -theory
(cohomology for compact spaces) (homology for C^* -algebras) (homology for rings)

What is a homology for C^* -algebras?

First, recall that we say a sequence of objects and morphisms

$$\dots \longrightarrow A \xrightarrow{f} B \xrightarrow{g} C \longrightarrow \dots$$

is **exact at B** if $\text{im } f = \ker g$. We say a sequence is **exact** if it is exact at all locations.

What is a homology for C^* -algebras?

First, recall that we say a sequence of objects and morphisms

$$\dots \longrightarrow A \xrightarrow{f} B \xrightarrow{g} C \longrightarrow \dots$$

is **exact at B** if $\text{im } f = \ker g$. We say a sequence is **exact** if it is exact at all locations.

A **short exact sequence** is an exact sequence of the form

$$0 \longrightarrow A \xrightarrow{f} B \xrightarrow{g} C \longrightarrow 0.$$

Note that if A , B , and C are C^* -algebras, then $\text{im } f = \ker g$, f is injective, g is surjective, A may be identified with an ideal in B , and $C \cong B/A$. So essentially any short exact sequence looks like

$$0 \longrightarrow I \xrightarrow{i} A \xrightarrow{\pi} A/I \longrightarrow 0.$$

for a C^* -algebra A and an ideal I of A .

What is a homology for C^* -algebras?

Motivation: Algebraic Topology

What is a homology for C^* -algebras?

Motivation: Algebraic Topology

To begin, a **homology** consists of a sequence of covariant functors $H_n : \mathbf{C}^* \rightarrow \mathbf{AbGp}$ for each $n \in \mathbb{N} \cup \{0\}$.

Notation for the functor H_n :

$$\begin{aligned} A &\rightsquigarrow H_n(A) \\ f : A \rightarrow B &\rightsquigarrow f_n : H_n(A) \rightarrow H_n(B) \end{aligned}$$

What is a homology for C^* -algebras?

Motivation: Algebraic Topology

To begin, a **homology** consists of a sequence of covariant functors $H_n : \mathbf{C}^* \rightarrow \mathbf{AbGp}$ for each $n \in \mathbb{N} \cup \{0\}$.

Notation for the functor H_n :

$$\begin{aligned} A &\rightsquigarrow H_n(A) \\ f : A \rightarrow B &\rightsquigarrow f_n : H_n(A) \rightarrow H_n(B) \end{aligned}$$

We require each H_n functor to be **half-exact**: For each $n \in \mathbb{N} \cup \{0\}$, whenever we have a short exact sequence

$$0 \longrightarrow A \xrightarrow{f} B \xrightarrow{g} C \longrightarrow 0$$

we may apply H_n to get a sequence

$$H_n(A) \xrightarrow{f_n} H_n(B) \xrightarrow{g_n} H_n(C)$$

that is exact at $H_n(B)$. (But typically not at $H_n(A)$ or $H_n(C)$.)

What is a homology for C^* -algebras?

Thus, when we have a short exact sequence

$$0 \longrightarrow A \xrightarrow{f} B \xrightarrow{g} C \longrightarrow 0$$

we may apply each H_n to get

$$H_0(A) \xrightarrow{f_0} H_0(B) \xrightarrow{g_0} H_0(C)$$

$$H_1(A) \xrightarrow{f_1} H_1(B) \xrightarrow{g_1} H_1(C)$$

$$H_2(A) \xrightarrow{f_2} H_2(B) \xrightarrow{g_2} H_2(C)$$

$$\vdots \qquad \qquad \qquad \vdots$$

For each n we require a **connecting homomorphism** $\delta_n : H_n(C) \rightarrow H_{n+1}(A)$ that makes a long exact sequence when inserted above. That is . . .

What is a homology for C^* -algebras?

$$\begin{array}{ccccc} H_0(A) & \xrightarrow{f_0} & H_0(B) & \xrightarrow{g_0} & H_0(C) \\ & & & \searrow \delta_0 & \\ H_1(A) & \xleftarrow{f_1} & H_1(B) & \xrightarrow{g_1} & H_1(C) \\ & & & \searrow \delta_1 & \\ H_2(A) & \xleftarrow{f_2} & H_2(B) & \xrightarrow{g_2} & H_2(C) \\ & & \vdots & & \vdots \\ & \swarrow & & & \\ & & \vdots & & \vdots \end{array}$$

We usually write this long exact sequence horizontally.

$$H_0(A) \xrightarrow{f_0} H_0(B) \xrightarrow{g_0} H_0(C) \xrightarrow{\delta_0} H_1(A) \xleftarrow{f_1} H_1(B) \xrightarrow{g_1} H_1(C) \xrightarrow{\delta_1} \dots$$

What is a homology for C^* -algebras?

In topology (when we assign long exact sequences of abelian groups to topological spaces), one can build the H_n -groups in different ways.

However, there is an axiomatization of a “unique” homology. One can prove that if the Eilenberg-Steenrod Axioms are satisfied, then the H_n -groups you get are the same (at least, on large classes of spaces).

What is a homology for C^* -algebras?

In topology (when we assign long exact sequences of abelian groups to topological spaces), one can build the H_n -groups in different ways.

However, there is an axiomatization of a “unique” homology. One can prove that if the Eilenberg-Steenrod Axioms are satisfied, then the H_n -groups you get are the same (at least, on large classes of spaces).

In a [cohomology](#) one uses contravariant functors, and you “reverse the arrows”.

What is a homology for C^* -algebras?

In topology (when we assign long exact sequences of abelian groups to topological spaces), one can build the H_n -groups in different ways.

However, there is an axiomatization of a “unique” homology. One can prove that if the Eilenberg-Steenrod Axioms are satisfied, then the H_n -groups you get are the same (at least, on large classes of spaces).

In a **cohomology** one uses contravariant functors, and you “reverse the arrows”.

Our homology for C^* -algebras is called **K -theory** and we’ll use the symbol K_n , in place of H_n , for our functors.

How do we build/define our K_n -groups? We look to topological K -theory, which was developed first, for motivation and inspiration.

Motivation: Topological K -theory

Topological K -theory is a cohomology for compact Hausdorff spaces.

The Big Idea: Fix a compact Hausdorff space X . The 0^{th} K -group for X is constructed using vector bundles over X , and the other groups are obtained by “suspending”; i.e., the n^{th} group is the 0^{th} group of the n^{th} suspension $S^n X$.

Motivation: Topological K -theory

Topological K -theory is a cohomology for compact Hausdorff spaces.

The Big Idea: Fix a compact Hausdorff space X . The 0^{th} K -group for X is constructed using vector bundles over X , and the other groups are obtained by “suspending”; i.e., the n^{th} group is the 0^{th} group of the n^{th} suspension $S^n X$.

How do we generalize to C^* -algebras (and rings)?

Noncommutative topology: We use the following functor

$$\begin{aligned} X &\rightsquigarrow C(X) \\ f : X \rightarrow Y &\rightsquigarrow f^* : C(Y) \rightarrow C(X) \\ &\text{where } f^*(g) := g \circ f \end{aligned}$$

Note: This functor is contravariant.

Motivation: Topological K -theory

Swan's Theorem: The category of vector bundles over a compact space X is equivalent (i.e., isomorphic in the category sense) to the category of finitely-generated projective modules over $C(X)$.

Finitely-generated: has a finite spanning set.

Projective: A module P is *projective* if for every surjective module homomorphism $f : N \rightarrow M$ and every module homomorphism $g : P \rightarrow M$, there exists a module homomorphism $h : P \rightarrow N$ such that $f \circ h = g$.

$$\begin{array}{ccc} & N & \\ \exists h \nearrow & \downarrow f & \\ P & \xrightarrow{g} & M \end{array}$$

(This is the definition of projective module, but it is equivalent to a handful of other properties.)

Motivation: Topological K -theory

Topological K -theory for a locally compact space X

0^{th} group formed using (isomorphism classes of) Vector Bundles over X .

Higher groups obtained by “suspending”.

Motivation: Topological K -theory

Topological K -theory for a locally compact space X

0^{th} group formed using (isomorphism classes of) Vector Bundles over X .
Higher groups obtained by “suspending”.

Operator (resp. Algebraic) K -theory for a C^* -algebra (resp. ring) R

0^{th} group formed using (isomorphism classes) of Finitely-Generated Projective Modules over R .

Higher groups obtained by “suspending”.

Motivation: Topological K -theory

Let R be a C^* -algebra, and let M be a projective module over R .

Motivation: Topological K -theory

Let R be a C^* -algebra, and let M be a projective module over R .

Then M is a direct summand of a free module; i.e., there exists N such that $M \oplus N$ is free.

Motivation: Topological K -theory

Let R be a C^* -algebra, and let M be a projective module over R .

Then M is a direct summand of a free module; i.e., there exists N such that $M \oplus N$ is free. If M is finitely generated, this free module can be chosen of finite rank; i.e., there exists $n \in \mathbb{N}$ such that

$$M \oplus N \cong R^n.$$

Motivation: Topological K -theory

Let R be a C^* -algebra, and let M be a projective module over R .

Then M is a direct summand of a free module; i.e., there exists N such that $M \oplus N$ is free. If M is finitely generated, this free module can be chosen of finite rank; i.e., there exists $n \in \mathbb{N}$ such that

$$M \oplus N \cong R^n.$$

This means M is a subspace of R^n . But, as you know, $\text{End } R^n \cong M_n(R)$, and we can identify the subspace M with the image of the projection $p \in M_n(R)$ onto M .

Motivation: Topological K -theory

Let R be a C^* -algebra, and let M be a projective module over R .

Then M is a direct summand of a free module; i.e., there exists N such that $M \oplus N$ is free. If M is finitely generated, this free module can be chosen of finite rank; i.e., there exists $n \in \mathbb{N}$ such that

$$M \oplus N \cong R^n.$$

This means M is a subspace of R^n . But, as you know, $\text{End } R^n \cong M_n(R)$, and we can identify the subspace M with the image of the projection $p \in M_n(R)$ onto M .

Q: When will two subspaces of R^n be isomorphic?

A: When there is an isomorphism (i.e., a partial isometry) between them. If p and q are the associated projections, this occurs iff there exists $v \in M_n(R)$ with $p = vv^*$ and $q = v^*v$. Murray-von Neumann equivalence!

Motivation: Topological K -theory

Topological K -theory for a locally compact space X

0^{th} group formed using (isomorphism classes of) Vector Bundles over X .
Higher groups obtained by “suspending”.

Operator (resp. Algebraic) K -theory for a C^* -algebra (resp. ring) R

0^{th} group formed using (isomorphism classes) of Finitely-Generated Projective Modules over R .

Motivation: Topological K -theory

Topological K -theory for a locally compact space X

0^{th} group formed using (isomorphism classes of) Vector Bundles over X .
Higher groups obtained by “suspending”.

Operator (resp. Algebraic) K -theory for a C^* -algebra (resp. ring) R

0^{th} group formed using (isomorphism classes) of Finitely-Generated Projective Modules over R .

. . . or equivalently . . .

0^{th} group constructed using Murray-von Neumann equivalence classes of projections (resp. idempotents) in square matrices over the C^* -algebra (resp. ring).

Higher groups obtained by “suspending”.

Motivation: Topological K -theory

Topological K -theory for a locally compact space X

0^{th} group formed using (isomorphism classes of) Vector Bundles over X .
Higher groups obtained by “suspending”.

Operator (resp. Algebraic) K -theory for a C^* -algebra (resp. ring) R

0^{th} group formed using (isomorphism classes) of Finitely-Generated Projective Modules over R .

. . . or equivalently . . .

0^{th} group constructed using Murray-von Neumann equivalence classes of projections (resp. idempotents) in square matrices over the C^* -algebra (resp. ring).

Higher groups obtained by “suspending”.

Let's focus on constructing K_0 for C^* -algebras and go through details.

Constructing the K_0 -group

Let A be a C^* -algebra. If p and q are projections in A , then $p + q$ may not be a projection. (It is precisely when $p \perp q$.)

Constructing the K_0 -group

Let A be a C^* -algebra. If p and q are projections in A , then $p + q$ may not be a projection. (It is precisely when $p \perp q$.)

However, in $M_2(A)$ we can identify p with $\begin{pmatrix} p & 0 \\ 0 & 0 \end{pmatrix}$, and we can identify q with $\begin{pmatrix} q & 0 \\ 0 & 0 \end{pmatrix} \sim \begin{pmatrix} 0 & 0 \\ 0 & q \end{pmatrix}$.

We can then define a sum

$$p \oplus q := \begin{pmatrix} p & 0 \\ 0 & q \end{pmatrix}.$$

Constructing the K_0 -group

Let A be a C^* -algebra. If p and q are projections in A , then $p + q$ may not be a projection. (It is precisely when $p \perp q$.)

However, in $M_2(A)$ we can identify p with $\begin{pmatrix} p & 0 \\ 0 & 0 \end{pmatrix}$, and we can identify q with $\begin{pmatrix} q & 0 \\ 0 & 0 \end{pmatrix} \sim \begin{pmatrix} 0 & 0 \\ 0 & q \end{pmatrix}$.

We can then define a sum

$$p \oplus q := \begin{pmatrix} p & 0 \\ 0 & q \end{pmatrix}.$$

Likewise for $p \in M_n(A)$ and $q \in M_k(A)$, we can define

$$p \oplus q := \begin{pmatrix} p & 0 \\ 0 & q \end{pmatrix} \in M_{n+k}(A).$$

The K_0 -group for Unital C^* -algebras

Let A be a unital C^* -algebra. Embed $M_n(A)$ in $M_{n+1}(A)$ by $x \mapsto \begin{pmatrix} x & 0 \\ 0 & 0 \end{pmatrix}$. Define

$$M_\infty(A) := \bigcup_{n=1}^{\infty} M_n(A).$$

Note: $M_\infty(A)$ is the non-closed $*$ -algebra of infinite matrices that have only finitely many nonzero entries. (Also, $\overline{M_\infty(\mathbb{C})} = \mathcal{K}(\mathcal{H})$.)

The K_0 -group for Unital C^* -algebras

Let A be a unital C^* -algebra. Embed $M_n(A)$ in $M_{n+1}(A)$ by $x \mapsto \begin{pmatrix} x & 0 \\ 0 & 0 \end{pmatrix}$.
Define

$$M_\infty(A) := \bigcup_{n=1}^{\infty} M_n(A).$$

Note: $M_\infty(A)$ is the non-closed $*$ -algebra of infinite matrices that have only finitely many nonzero entries. (Also, $\overline{M_\infty(\mathbb{C})} = \mathcal{K}(\mathcal{H})$.)

Define

$$V(A) := \{[p] : p \in \text{Proj } M_\infty(A)\}$$

with

$$[p] + [q] := \left[\begin{pmatrix} p & 0 \\ 0 & q \end{pmatrix} \right].$$

The K_0 -group for Unital C^* -algebras

Let A be a unital C^* -algebra. Embed $M_n(A)$ in $M_{n+1}(A)$ by $x \mapsto \begin{pmatrix} x & 0 \\ 0 & 0 \end{pmatrix}$.

Define

$$M_\infty(A) := \bigcup_{n=1}^{\infty} M_n(A).$$

Note: $M_\infty(A)$ is the non-closed $*$ -algebra of infinite matrices that have only finitely many nonzero entries. (Also, $\overline{M_\infty(\mathbb{C})} = \mathcal{K}(\mathcal{H})$.)

Define

$$V(A) := \{[p] : p \in \text{Proj } M_\infty(A)\}$$

with

$$[p] + [q] := \left[\begin{pmatrix} p & 0 \\ 0 & q \end{pmatrix} \right].$$

(The symbol V is a historical carryover — it stands for "vector bundle".)

Fact: $V(A)$ is an abelian semigroup with identity (i.e., an abelian monoid).

We want a group.

The Grothendieck Group of a Semigroup

Let $(V, +)$ be an abelian semigroup with identity.

The Grothendieck Group of a Semigroup

Let $(V, +)$ be an abelian semigroup with identity.

Consider a pair $(h, k) \in V \times V$ and “think of it” representing $h - k$.

Define an equivalence relation \equiv on $V \times V$ by

$$(h_1, k_1) \equiv (h_2, k_2) \iff \exists x \in V \text{ s.t. } h_1 + k_2 + x = h_2 + k_1 + x.$$

Why the x ? To get transitivity.

The Grothendieck Group of a Semigroup

Let $(V, +)$ be an abelian semigroup with identity.

Consider a pair $(h, k) \in V \times V$ and “think of it” representing $h - k$.

Define an equivalence relation \equiv on $V \times V$ by

$$(h_1, k_1) \equiv (h_2, k_2) \iff \exists x \in V \text{ s.t. } h_1 + k_2 + x = h_2 + k_1 + x.$$

Why the x ? To get transitivity.

The **Grothendieck Group** is the set of equivalence classes

$$\text{Groth } V := \{[(h, k)] : h, k \in V\} \text{ w/ } [(h_1, k_1)] + [(h_2, k_2)] = [(h_1 + h_2, k_1 + k_2)].$$

The Grothendieck Group of a Semigroup

Let $(V, +)$ be an abelian semigroup with identity.

Consider a pair $(h, k) \in V \times V$ and “think of it” representing $h - k$.

Define an equivalence relation \equiv on $V \times V$ by

$$(h_1, k_1) \equiv (h_2, k_2) \iff \exists x \in V \text{ s.t. } h_1 + k_2 + x = h_2 + k_1 + x.$$

Why the x ? To get transitivity.

The **Grothendieck Group** is the set of equivalence classes

$$\text{Groth } V := \{[(h, k)] : h, k \in V\} \text{ w/ } [(h_1, k_1)] + [(h_2, k_2)] = [(h_1 + h_2, k_1 + k_2)].$$

We often write $[(h, k)]$ as the formal difference $h - k$.

But keep in mind: $h_1 - k_1 = h_2 - k_2$ iff $\exists x$ s.t. $h_1 + k_2 + x = h_2 + k_1 + x$.

$\text{Groth } V$ is an abelian group and universal for V in the following sense:

We can “include” $V \rightarrow \text{Groth } V$ by $h \mapsto (h, 0)$, (this isn’t always injective). If G is a group and there is a homomorphism $\phi : V \rightarrow G$, then ϕ extends to $\tilde{\phi} : \text{Groth } V \rightarrow G$ by $\tilde{\phi}(h - k) = \phi(h) - \phi(k)$.

Examples:

Let $V = \{0, 1, 2, 3, \dots\}$ with $+$. Then $\text{Groth } V \cong \mathbb{Z}$.

Examples:

Let $V = \{0, 1, 2, 3, \dots\}$ with $+$. Then $\text{Groth } V \cong \mathbb{Z}$.

Let $V = \{0, 1, 2, 3, \dots\} \cup \{\infty\}$ with $+$. Then $\text{Groth } V \cong 0$.
(b/c $x + \infty = y + \infty$ for all x, y)

Examples:

Let $V = \{0, 1, 2, 3, \dots\}$ with $+$. Then $\text{Groth } V \cong \mathbb{Z}$.

Let $V = \{0, 1, 2, 3, \dots\} \cup \{\infty\}$ with $+$. Then $\text{Groth } V \cong 0$.
(b/c $x + \infty = y + \infty$ for all x, y)

Let $V = \{1, 2, 3, \dots\}$ with \times . Then $\text{Groth } V \cong \mathbb{Q}^+$.

Constructing the K_0 -group

Back to $K_0(A)$. . .

A is a unital C^* -algebra.

$$V(A) := \{[p] : p \in \text{Proj } M_\infty(A)\} \text{ with } [p] + [q] = \left[\begin{pmatrix} p & 0 \\ 0 & q \end{pmatrix} \right].$$

Constructing the K_0 -group

Back to $K_0(A)$. . .

A is a unital C^* -algebra.

$$V(A) := \{[p] : p \in \text{Proj } M_\infty(A)\} \text{ with } [p] + [q] = \left[\begin{pmatrix} p & 0 \\ 0 & q \end{pmatrix} \right].$$

We then define

$$K_0(A) := \text{Groth } V(A) = \{[p] - [q] : p, q \in \text{Proj } M_\infty(A)\}.$$

Constructing the K_0 -group

Back to $K_0(A)$. . .

A is a unital C^* -algebra.

$$V(A) := \{[p] : p \in \text{Proj } M_\infty(A)\} \text{ with } [p] + [q] = \left[\begin{pmatrix} p & 0 \\ 0 & q \end{pmatrix} \right].$$

We then define

$$K_0(A) := \text{Groth } V(A) = \{[p] - [q] : p, q \in \text{Proj } M_\infty(A)\}.$$

Also, we want K_0 to be a functor, so if $h : A \rightarrow B$ is a $*$ -homomorphism, we define $h_0 : K_0(A) \rightarrow K_0(B)$ by

$$h_0([p] - [q]) = [h(p)] - [h(q)].$$

Constructing the K_0 -group

What about when A is nonunital?

Constructing the K_0 -group

What about when A is nonunital? Let A be a nonunital C^* -algebra. Let A^1 be its (minimal) unitization. We have a short exact sequence

$$0 \longrightarrow A \xrightarrow{i} A^1 \xrightarrow{\pi} \mathbb{C} \longrightarrow 0$$

and both A^1 and \mathbb{C} are unital, so using our prior definition we obtain $\pi_0 : K_0(A^1) \rightarrow K_0(\mathbb{C}) \cong \mathbb{Z}$. We then define

$$K_0(A) := \ker \pi_0.$$

Constructing the K_0 -group

What about when A is nonunital? Let A be a nonunital C^* -algebra. Let A^1 be its (minimal) unitization. We have a short exact sequence

$$0 \longrightarrow A \xrightarrow{i} A^1 \xrightarrow{\pi} \mathbb{C} \longrightarrow 0$$

and both A^1 and \mathbb{C} are unital, so using our prior definition we obtain $\pi_0 : K_0(A^1) \rightarrow K_0(\mathbb{C}) \cong \mathbb{Z}$. We then define

$$K_0(A) := \ker \pi_0.$$

Fact: It turns out, that $K_0(A^1) \cong K_0(A) \oplus \mathbb{Z}$ when A is nonunital.

Fact: If A has a countable approximate unit consisting of projections, then

$$K_0(A) \cong \text{Groth } V(A) = \{[p] - [q] : p, q \in \text{Proj } M_\infty(A)\}.$$

Examples of K_0

\mathbb{C} , $M_n(\mathbb{C})$, and $\mathcal{K}(\mathcal{H})$

Projections in $M_\infty(\mathbb{C})$ are finite rank, so $V(\mathbb{C}) \cong \{0, 1, 2, \dots\}$ and

$$K_0(\mathbb{C}) \cong \mathbb{Z}.$$

Examples of K_0

\mathbb{C} , $M_n(\mathbb{C})$, and $\mathcal{K}(\mathcal{H})$

Projections in $M_\infty(\mathbb{C})$ are finite rank, so $V(\mathbb{C}) \cong \{0, 1, 2, \dots\}$ and

$$K_0(\mathbb{C}) \cong \mathbb{Z}.$$

Likewise, $M_\infty(M_n(\mathbb{C})) = M_\infty(\mathbb{C})$, and projections in $\mathcal{K}(\mathcal{H})$ and $M_\infty(\mathcal{K}(\mathcal{H}))$ are finite rank, so $V(M_n(\mathbb{C})) \cong V(\mathcal{K}(\mathcal{H})) \cong \{0, 1, 2, \dots\}$ and

$$K_0(M_n(\mathbb{C})) \cong \mathbb{Z} \quad \text{and} \quad K_0(\mathcal{K}(\mathcal{H})) \cong \mathbb{Z}.$$

Examples of K_0

\mathbb{C} , $M_n(\mathbb{C})$, and $\mathcal{K}(\mathcal{H})$

Projections in $M_\infty(\mathbb{C})$ are finite rank, so $V(\mathbb{C}) \cong \{0, 1, 2, \dots\}$ and

$$K_0(\mathbb{C}) \cong \mathbb{Z}.$$

Likewise, $M_\infty(M_n(\mathbb{C})) = M_\infty(\mathbb{C})$, and projections in $\mathcal{K}(\mathcal{H})$ and $M_\infty(\mathcal{K}(\mathcal{H}))$ are finite rank, so $V(M_n(\mathbb{C})) \cong V(\mathcal{K}(\mathcal{H})) \cong \{0, 1, 2, \dots\}$ and

$$K_0(M_n(\mathbb{C})) \cong \mathbb{Z} \quad \text{and} \quad K_0(\mathcal{K}(\mathcal{H})) \cong \mathbb{Z}.$$

$B(\mathcal{H})$

In $M_\infty(B(\mathcal{H})) \cong B(\mathcal{H})$ all projections are either finite rank or have countably infinite rank. So $V(B(\mathcal{H})) \cong \{0, 1, 2, \dots\} \cup \{\infty\}$ and

$$K_0(B(\mathcal{H})) \cong \{0\}.$$

Examples of K_0

\mathbb{C} , $M_n(\mathbb{C})$, and $\mathcal{K}(\mathcal{H})$

Projections in $M_\infty(\mathbb{C})$ are finite rank, so $V(\mathbb{C}) \cong \{0, 1, 2, \dots\}$ and

$$K_0(\mathbb{C}) \cong \mathbb{Z}.$$

Likewise, $M_\infty(M_n(\mathbb{C})) = M_\infty(\mathbb{C})$, and projections in $\mathcal{K}(\mathcal{H})$ and $M_\infty(\mathcal{K}(\mathcal{H}))$ are finite rank, so $V(M_n(\mathbb{C})) \cong V(\mathcal{K}(\mathcal{H})) \cong \{0, 1, 2, \dots\}$ and

$$K_0(M_n(\mathbb{C})) \cong \mathbb{Z} \quad \text{and} \quad K_0(\mathcal{K}(\mathcal{H})) \cong \mathbb{Z}.$$

$B(\mathcal{H})$

In $M_\infty(B(\mathcal{H})) \cong B(\mathcal{H})$ all projections are either finite rank or have countably infinite rank. So $V(B(\mathcal{H})) \cong \{0, 1, 2, \dots\} \cup \{\infty\}$ and

$$K_0(B(\mathcal{H})) \cong \{0\}.$$

$\mathcal{C}(\mathcal{H}) := B(\mathcal{H})/\mathcal{K}(\mathcal{H})$

In $\mathcal{C}(\mathcal{H})$ and $M_\infty(\mathcal{C}(\mathcal{H}))$ all finite-rank projections are equivalent, so $V(\mathcal{C}(\mathcal{H})) = \{0, \infty\}$ and

$$K_0(\mathcal{C}(\mathcal{H})) \cong \{0\}.$$

A Note on Equivalence in the K_0 -group

Let p and q be projections in A . We say p and q are . . .

Murray-von Neumann equivalent, denoted $p \sim q$ if there exists $v \in A$ with $p = vv^*$ and $q = v^*v$.

unitarily equivalent, denoted $p \sim_u q$, if there exists unitary $u \in A^1$ with $p = u^*qu$.

homotopic, denoted $p \sim_h q$, when p and q are connected by a norm-continuous path of projections in A .

A Note on Equivalence in the K_0 -group

Let p and q be projections in A . We say p and q are . . .

Murray-von Neumann equivalent, denoted $p \sim q$ if there exists $v \in A$ with $p = vv^*$ and $q = v^*v$.

unitarily equivalent, denoted $p \sim_u q$, if there exists unitary $u \in A^1$ with $p = u^*qu$.

homotopic, denoted $p \sim_h q$, when p and q are connected by a norm-continuous path of projections in A .

Facts:

$$p \sim_h q \implies p \sim_u q \implies p \sim q$$

$$p \sim q \implies \begin{pmatrix} p & 0 \\ 0 & 0 \end{pmatrix} \sim_u \begin{pmatrix} q & 0 \\ 0 & 0 \end{pmatrix} \quad \text{and} \quad p \sim_u q \implies \begin{pmatrix} p & 0 \\ 0 & 0 \end{pmatrix} \sim_h \begin{pmatrix} q & 0 \\ 0 & 0 \end{pmatrix}$$

A Note on Equivalence in the K_0 -group

Let p and q be projections in A . We say p and q are . . .

Murray-von Neumann equivalent, denoted $p \sim q$ if there exists $v \in A$ with $p = vv^*$ and $q = v^*v$.

unitarily equivalent, denoted $p \sim_u q$, if there exists unitary $u \in A^1$ with $p = u^*qu$.

homotopic, denoted $p \sim_h q$, when p and q are connected by a norm-continuous path of projections in A .


Facts:

$$p \sim_h q \implies p \sim_u q \implies p \sim q$$
$$p \sim q \implies \begin{pmatrix} p & 0 \\ 0 & 0 \end{pmatrix} \sim_u \begin{pmatrix} q & 0 \\ 0 & 0 \end{pmatrix} \quad \text{and} \quad p \sim_u q \implies \begin{pmatrix} p & 0 \\ 0 & 0 \end{pmatrix} \sim_h \begin{pmatrix} q & 0 \\ 0 & 0 \end{pmatrix}$$

So in $M_\infty(A)$ (and hence in $K_0(A)$) the Murray-von Neumann equivalence classes, unitary equivalence classes, and homotopy equivalence classes coincide.


The Higher K -groups

In topology, the **suspension** of a topological space X is intuitively obtained by stretching X into a cylinder and then collapsing both end faces to points. One views X as “suspended” between these end points.



The Higher K -groups

In topology, the **suspension** of a topological space X is intuitively obtained by stretching X into a cylinder and then collapsing both end faces to points. One views X as “suspended” between these end points.



The noncommutative version: If A is a C^* -algebra,

$$SA := \{f \in C([0, 1], A) : f(0) = f(1) = 0\}.$$

Equivalent descriptions:

$$SA \cong C_0((0, 1), A) \cong C_0(\mathbb{R}, A) \cong \{f \in C(\mathbb{T}, A) : f(1) = 0\}$$

The Higher K -groups

Higher K -groups are defined inductively. Given $K_0(A)$, we define

$$K_{n+1}(A) := K_n(SA) \quad \text{for } n = 0, 1, 2, \dots$$

So inductively we obtain $K_n(A) := K_0(S^n A)$.

The Higher K -groups

Higher K -groups are defined inductively. Given $K_0(A)$, we define

$$K_{n+1}(A) := K_n(SA) \quad \text{for } n = 0, 1, 2, \dots$$

So inductively we obtain $K_n(A) := K_0(S^n A)$.

Although the K_1 -group is defined as $K_1(A) := K_0(SA)$, we can also obtain a description in terms of unitaries . . .

The K_1 -group

Define

$$A^+ := \begin{cases} A^1 & \text{if } A \text{ is nonunital} \\ A & \text{if } A \text{ is unital.} \end{cases}$$

The K_1 -group

Define

$$A^+ := \begin{cases} A^1 & \text{if } A \text{ is nonunital} \\ A & \text{if } A \text{ is unital.} \end{cases}$$

Let $U_n(A^+)$ denote set of unitaries in $M_n(A^+)$. We can embed $U_n(A^+)$ in $U_{n+1}(A^+)$ by $x \mapsto \begin{pmatrix} x & 0 \\ 0 & 1 \end{pmatrix}$, and we define $U_\infty(A^+) := \bigcup_{n=1}^\infty U_n(A^+)$.

The K_1 -group

Define

$$A^+ := \begin{cases} A^1 & \text{if } A \text{ is nonunital} \\ A & \text{if } A \text{ is unital.} \end{cases}$$

Let $U_n(A^+)$ denote set of unitaries in $M_n(A^+)$. We can embed $U_n(A^+)$ in $U_{n+1}(A^+)$ by $x \mapsto \begin{pmatrix} x & 0 \\ 0 & 1 \end{pmatrix}$, and we define $U_\infty(A^+) := \bigcup_{n=1}^\infty U_n(A^+)$.

We say $u, v \in U_n(A^+)$ are **homotopic** if there is a norm-continuous path $\gamma : [0, 1] \rightarrow U_n(A^+)$ with $\phi(0) = u$ and $\phi(1) = v$.

The K_1 -group

Define

$$A^+ := \begin{cases} A^1 & \text{if } A \text{ is nonunital} \\ A & \text{if } A \text{ is unital.} \end{cases}$$

Let $U_n(A^+)$ denote set of unitaries in $M_n(A^+)$. We can embed $U_n(A^+)$ in $U_{n+1}(A^+)$ by $x \mapsto \begin{pmatrix} x & 0 \\ 0 & 1 \end{pmatrix}$, and we define $U_\infty(A^+) := \bigcup_{n=1}^\infty U_n(A^+)$.

We say $u, v \in U_n(A^+)$ are **homotopic** if there is a norm-continuous path $\gamma : [0, 1] \rightarrow U_n(A^+)$ with $\phi(0) = u$ and $\phi(1) = v$.

Given $u, v \in U_\infty(A^+)$ with $u \in U_n(A^+)$ and $v \in U_m(A^+)$, we define $u \sim_h v$ if $\exists k \geq \max\{m, n\}$ s.t. $\begin{pmatrix} u & 0 \\ 0 & 1_{k-n} \end{pmatrix}$ and $\begin{pmatrix} v & 0 \\ 0 & 1_{k-m} \end{pmatrix}$ are homotopic.

The K_1 -group

Define

$$A^+ := \begin{cases} A^1 & \text{if } A \text{ is nonunital} \\ A & \text{if } A \text{ is unital.} \end{cases}$$

Let $U_n(A^+)$ denote set of unitaries in $M_n(A^+)$. We can embed $U_n(A^+)$ in $U_{n+1}(A^+)$ by $x \mapsto \begin{pmatrix} x & 0 \\ 0 & 1 \end{pmatrix}$, and we define $U_\infty(A^+) := \bigcup_{n=1}^\infty U_n(A^+)$.

We say $u, v \in U_n(A^+)$ are **homotopic** if there is a norm-continuous path $\gamma : [0, 1] \rightarrow U_n(A^+)$ with $\phi(0) = u$ and $\phi(1) = v$.

Given $u, v \in U_\infty(A^+)$ with $u \in U_n(A^+)$ and $v \in U_m(A^+)$, we define $u \sim_h v$ if $\exists k \geq \max\{m, n\}$ s.t. $\begin{pmatrix} u & 0 \\ 0 & 1_{k-n} \end{pmatrix}$ and $\begin{pmatrix} v & 0 \\ 0 & 1_{k-m} \end{pmatrix}$ are homotopic.

We define

$$K_1(A) := U_\infty(A^+) / \sim_h \quad \text{with} \quad [u]_h + [v]_h := \left[\begin{pmatrix} u & 0 \\ 0 & v \end{pmatrix} \right]_h$$

The K_1 -group

Define

$$A^+ := \begin{cases} A^1 & \text{if } A \text{ is nonunital} \\ A & \text{if } A \text{ is unital.} \end{cases}$$

Let $U_n(A^+)$ denote set of unitaries in $M_n(A^+)$. We can embed $U_n(A^+)$ in $U_{n+1}(A^+)$ by $x \mapsto \begin{pmatrix} x & 0 \\ 0 & 1 \end{pmatrix}$, and we define $U_\infty(A^+) := \bigcup_{n=1}^\infty U_n(A^+)$.

We say $u, v \in U_n(A^+)$ are **homotopic** if there is a norm-continuous path $\gamma : [0, 1] \rightarrow U_n(A^+)$ with $\phi(0) = u$ and $\phi(1) = v$.

Given $u, v \in U_\infty(A^+)$ with $u \in U_n(A^+)$ and $v \in U_m(A^+)$, we define $u \sim_h v$ if $\exists k \geq \max\{m, n\}$ s.t. $\begin{pmatrix} u & 0 \\ 0 & 1_{k-n} \end{pmatrix}$ and $\begin{pmatrix} v & 0 \\ 0 & 1_{k-m} \end{pmatrix}$ are homotopic.

We define

$$K_1(A) := U_\infty(A^+) / \sim_h \quad \text{with} \quad [u]_h + [v]_h := \left[\begin{pmatrix} u & 0 \\ 0 & v \end{pmatrix} \right]_h$$

Fact: $K_1(A)$ is an abelian group; moreover $-[u]_h = [u^*]_h$.

K_1 is a functor: If $\phi : A \rightarrow B$, it extends to $\tilde{\phi} : M_\infty(A^+) \rightarrow M_\infty(B^+)$ and we define $\phi_1 : K_1(A) \rightarrow K_1(B)$ by $\phi_1([u]_h) = [\tilde{\phi}(u)]_h$

Examples of K_1

The K_1 -group is a bit harder to compute at this stage. But with some work, one can prove that all unitaries in $U_\infty(\mathbb{C})$ and $U_\infty(B(\mathcal{H}))$ are homotopic, giving

$$K_1(\mathbb{C}) \cong K_1(M_n(\mathbb{C})) \cong K_1(\mathcal{K}(\mathcal{H})) \cong K_1(B(\mathcal{H})) \cong \{0\}.$$

Examples of K_1

The K_1 -group is a bit harder to compute at this stage. But with some work, one can prove that all unitaries in $U_\infty(\mathbb{C})$ and $U_\infty(B(\mathcal{H}))$ are homotopic, giving

$$K_1(\mathbb{C}) \cong K_1(M_n(\mathbb{C})) \cong K_1(\mathcal{K}(\mathcal{H})) \cong K_1(B(\mathcal{H})) \cong \{0\}.$$

We'll show some tricks for computing more K_1 -groups later.

The Index Maps

At this point we have our functors K_n , but to obtain a homology we also need **connecting maps** (sometimes called **index maps**); i.e., for each C^* -algebra A and each ideal I of A , we need to construct a map

$$\delta_n : K_n(A/I) \rightarrow K_{n+1}(I) \quad \text{for each } n = 0, 1, \dots$$

The Index Maps

At this point we have our functors K_n , but to obtain a homology we also need **connecting maps** (sometimes called **index maps**); i.e., for each C^* -algebra A and each ideal I of A , we need to construct a map

$$\delta_n : K_n(A/I) \rightarrow K_{n+1}(I) \quad \text{for each } n = 0, 1, \dots$$

I'll spare you the details, but the index maps do exist. Moreover, it can be proven that each is unique up to sign, so despite what may seem to be a complicated or unmotivated construction, we are assured we have obtained the correct map in the end.

The Index Maps

At this point we have our functors K_n , but to obtain a homology we also need **connecting maps** (sometimes called **index maps**); i.e., for each C^* -algebra A and each ideal I of A , we need to construct a map

$$\delta_n : K_n(A/I) \rightarrow K_{n+1}(I) \quad \text{for each } n = 0, 1, \dots$$

I'll spare you the details, but the index maps do exist. Moreover, it can be proven that each is unique up to sign, so despite what may seem to be a complicated or unmotivated construction, we are assured we have obtained the correct map in the end.

Thus for any ideal I in A , we map apply K -theory to the short exact sequence $0 \rightarrow I \rightarrow A \rightarrow A/I \rightarrow 0$ to obtain a long exact sequence

$$K_0(I) \longrightarrow K_0(A) \longrightarrow K_0(A/I) \xrightarrow{\delta_0} K_1(I) \longrightarrow K_1(A) \longrightarrow K_1(A/I) \xrightarrow{\delta_1} \dots$$

The Index Maps

At this point we have our functors K_n , but to obtain a homology we also need **connecting maps** (sometimes called **index maps**); i.e., for each C^* -algebra A and each ideal I of A , we need to construct a map

$$\delta_n : K_n(A/I) \rightarrow K_{n+1}(I) \quad \text{for each } n = 0, 1, \dots$$

I'll spare you the details, but the index maps do exist. Moreover, it can be proven that each is unique up to sign, so despite what may seem to be a complicated or unmotivated construction, we are assured we have obtained the correct map in the end.

Thus for any ideal I in A , we map apply K -theory to the short exact sequence $0 \rightarrow I \rightarrow A \rightarrow A/I \rightarrow 0$ to obtain a long exact sequence

$$K_0(I) \longrightarrow K_0(A) \longrightarrow K_0(A/I) \xrightarrow{\delta_0} K_1(I) \longrightarrow K_1(A) \longrightarrow K_1(A/I) \xrightarrow{\delta_1} \dots$$

In addition, a truly remarkable fact emerges during the construction of the index maps . . .

Bott Periodicity

It turns out that $K_0(A) \cong K_2(A)$ for any C^* -algebra A . (Wow!)

Bott Periodicity

It turns out that $K_0(A) \cong K_2(A)$ for any C^* -algebra A . (Wow!)

This implies all the higher K -groups after K_1 are redundant. For instance,

$$K_3(A) := K_2(SA) \cong K_0(SA) = K_1(A).$$

Bott Periodicity

It turns out that $K_0(A) \cong K_2(A)$ for any C^* -algebra A . (Wow!)

This implies all the higher K -groups after K_1 are redundant. For instance,

$$K_3(A) := K_2(SA) \cong K_0(SA) = K_1(A).$$

Inductively, we obtain

$$K_0(A) \cong K_2(A) \cong K_4(A) \cong K_6(A) \cong \dots$$

and
$$K_1(A) \cong K_3(A) \cong K_5(A) \cong K_7(A) \cong \dots$$

Thus there are really only two distinct K -groups: $K_0(A)$ and $K_1(A)$.

Bott Periodicity

It turns out that $K_0(A) \cong K_2(A)$ for any C^* -algebra A . (Wow!)

This implies all the higher K -groups after K_1 are redundant. For instance,

$$K_3(A) := K_2(SA) \cong K_0(SA) = K_1(A).$$

Inductively, we obtain

$$K_0(A) \cong K_2(A) \cong K_4(A) \cong K_6(A) \cong \dots$$

and

$$K_1(A) \cong K_3(A) \cong K_5(A) \cong K_7(A) \cong \dots$$

Thus there are really only two distinct K -groups: $K_0(A)$ and $K_1(A)$.

Also, since the K_0 -group and the K_2 -group of any C^* -algebra agree, for any short exact sequence $0 \rightarrow I \rightarrow A \rightarrow A/I \rightarrow 0$, the corresponding long exact sequence

$$K_0(I) \longrightarrow K_0(A) \longrightarrow K_0(A/I) \xrightarrow{\delta_0} K_1(I) \longrightarrow K_1(A) \longrightarrow K_1(A/I) \xrightarrow{\delta_1} \dots$$

wraps around on itself . . .

Theorem (The Cyclic 6-term Exact Sequence)

For any C^* -algebra A and any ideal I of A , applying K -theory to the short exact sequence

$$0 \longrightarrow I \xrightarrow{i} A \xrightarrow{\pi} A/I \longrightarrow 0$$

yields the cyclic 6-term exact sequence

$$\begin{array}{ccccc} K_0(I) & \xrightarrow{i_0} & K_0(A) & \xrightarrow{\pi_0} & K_0(A/I) \\ \delta_1 \uparrow & & & & \downarrow \delta_0 \\ K_1(A/I) & \xleftarrow{\pi_1} & K_1(A) & \xleftarrow{i_1} & K_1(I) \end{array}$$

Topological K -theory also has Bott periodicity of period 2. Algebraic K -theory does not have Bott periodicity.

Fun Fact: If you work over \mathbb{R} instead of \mathbb{C} in Topological or Operator K -theory, you get period 8 and a cyclic 24-term exact sequence.

The 6-term exact sequence can be useful for computing K -groups.

The 6-term exact sequence can be useful for computing K -groups.

Example: We know the K -groups for $\mathcal{K}(\mathcal{H})$ and $B(\mathcal{H})$. We can use them to calculate the K -groups of the Calkin algebra $\mathcal{C}(\mathcal{H}) := B(\mathcal{H})/\mathcal{K}(\mathcal{H})$.

The 6-term exact sequence can be useful for computing K -groups.

Example: We know the K -groups for $\mathcal{K}(\mathcal{H})$ and $B(\mathcal{H})$. We can use them to calculate the K -groups of the Calkin algebra $\mathcal{C}(\mathcal{H}) := B(\mathcal{H})/\mathcal{K}(\mathcal{H})$. Applying K -theory to $0 \rightarrow \mathcal{K}(\mathcal{H}) \rightarrow B(\mathcal{H}) \rightarrow \mathcal{C}(\mathcal{H}) \rightarrow 0$ we get

$$\begin{array}{ccccc} K_0(\mathcal{K}(\mathcal{H})) & \longrightarrow & K_0(B(\mathcal{H})) & \longrightarrow & K_0(\mathcal{C}(\mathcal{H})) \\ & & & & \downarrow \\ & \uparrow & & & \\ K_1(\mathcal{C}(\mathcal{H})) & \longleftarrow & K_1(B(\mathcal{H})) & \longleftarrow & K_1(\mathcal{K}(\mathcal{H})) \end{array}$$

The 6-term exact sequence can be useful for computing K -groups.

Example: We know the K -groups for $\mathcal{K}(\mathcal{H})$ and $B(\mathcal{H})$. We can use them to calculate the K -groups of the Calkin algebra $\mathcal{C}(\mathcal{H}) := B(\mathcal{H})/\mathcal{K}(\mathcal{H})$. Applying K -theory to $0 \rightarrow \mathcal{K}(\mathcal{H}) \rightarrow B(\mathcal{H}) \rightarrow \mathcal{C}(\mathcal{H}) \rightarrow 0$ we get

$$\begin{array}{ccccc}
 K_0(\mathcal{K}(\mathcal{H})) & \longrightarrow & K_0(B(\mathcal{H})) & \longrightarrow & K_0(\mathcal{C}(\mathcal{H})) \\
 \uparrow & & & & \downarrow \\
 K_1(\mathcal{C}(\mathcal{H})) & \longleftarrow & K_1(B(\mathcal{H})) & \longleftarrow & K_1(\mathcal{K}(\mathcal{H}))
 \end{array}$$

Substituting known values yields

$$\begin{array}{ccccc}
 \mathbb{Z} & \longrightarrow & 0 & \longrightarrow & K_0(\mathcal{C}(\mathcal{H})) \\
 \uparrow & & & & \downarrow \\
 K_1(\mathcal{C}(\mathcal{H})) & \longleftarrow & 0 & \longleftarrow & 0
 \end{array}$$

The 6-term exact sequence can be useful for computing K -groups.

Example: We know the K -groups for $\mathcal{K}(\mathcal{H})$ and $B(\mathcal{H})$. We can use them to calculate the K -groups of the Calkin algebra $\mathcal{C}(\mathcal{H}) := B(\mathcal{H})/\mathcal{K}(\mathcal{H})$. Applying K -theory to $0 \rightarrow \mathcal{K}(\mathcal{H}) \rightarrow B(\mathcal{H}) \rightarrow \mathcal{C}(\mathcal{H}) \rightarrow 0$ we get

$$\begin{array}{ccccc}
 K_0(\mathcal{K}(\mathcal{H})) & \longrightarrow & K_0(B(\mathcal{H})) & \longrightarrow & K_0(\mathcal{C}(\mathcal{H})) \\
 \uparrow & & & & \downarrow \\
 K_1(\mathcal{C}(\mathcal{H})) & \longleftarrow & K_1(B(\mathcal{H})) & \longleftarrow & K_1(\mathcal{K}(\mathcal{H}))
 \end{array}$$

Substituting known values yields

$$\begin{array}{ccccc}
 \mathbb{Z} & \longrightarrow & 0 & \longrightarrow & K_0(\mathcal{C}(\mathcal{H})) \\
 \uparrow & & & & \downarrow \\
 K_1(\mathcal{C}(\mathcal{H})) & \longleftarrow & 0 & \longleftarrow & 0
 \end{array}$$

So $K_1(\mathcal{C}(\mathcal{H})) \cong \mathbb{Z}$ and $K_0(\mathcal{C}(\mathcal{H})) \cong \{0\}$.

A covariant functor F from \mathbf{C}^* to \mathbf{AbGp} is . . .

- **Half Exact** when every short exact sequence $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$ is taken to an exact sequence $FA \rightarrow FB \rightarrow FC$.
- **Homotopy Invariant** If $\alpha : A \rightarrow B$ and $\beta : A \rightarrow B$ are homotopic (i.e., there exists a path of morphisms $\gamma_t : A \rightarrow B$, $t \in [0, 1]$ such that $t \mapsto \gamma_t(a)$ is norm continuous for all $a \in A$ and with $\gamma_0 = \alpha$ and $\gamma_1 = \beta$), then $\alpha_* = \beta_*$.
- **Stable** For any C^* -algebra A and any rank 1 projection $p \in \mathcal{K}(\mathcal{H})$, the morphism $a \mapsto a \otimes p$ from A to $A \otimes \mathcal{K}(\mathcal{H})$ induces an isomorphism from $F(A)$ onto $F(A \otimes \mathcal{K}(\mathcal{H}))$.
- **Continuous** if whenever $\{A_n, \phi_n\}_{n=1}^\infty$ is a countable directed sequence, then $F(\varinjlim(A_n, \phi_n)) = \varinjlim(F(A_n), \phi_{n*})$

A covariant functor F from \mathbf{C}^* to \mathbf{AbGp} is . . .

- **Half Exact** when every short exact sequence $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$ is taken to an exact sequence $FA \rightarrow FB \rightarrow FC$.
- **Homotopy Invariant** If $\alpha : A \rightarrow B$ and $\beta : A \rightarrow B$ are homotopic (i.e., there exists a path of morphisms $\gamma_t : A \rightarrow B$, $t \in [0, 1]$ such that $t \mapsto \gamma_t(a)$ is norm continuous for all $a \in A$ and with $\gamma_0 = \alpha$ and $\gamma_1 = \beta$), then $\alpha_* = \beta_*$.
- **Stable** For any C^* -algebra A and any rank 1 projection $p \in \mathcal{K}(\mathcal{H})$, the morphism $a \mapsto a \otimes p$ from A to $A \otimes \mathcal{K}(\mathcal{H})$ induces an isomorphism from $F(A)$ onto $F(A \otimes \mathcal{K}(\mathcal{H}))$.
- **Continuous** if whenever $\{A_n, \phi_n\}_{n=1}^\infty$ is a countable directed sequence, then $F(\varinjlim(A_n, \phi_n)) = \varinjlim(F(A_n), \phi_{n*})$

K_0 and K_1 are half exact, homotopy invariant, stable, and continuous.

A covariant functor F from \mathbf{C}^* to \mathbf{AbGp} is . . .

- **Half Exact** when every short exact sequence $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$ is taken to an exact sequence $FA \rightarrow FB \rightarrow FC$.
- **Homotopy Invariant** If $\alpha : A \rightarrow B$ and $\beta : A \rightarrow B$ are homotopic (i.e., there exists a path of morphisms $\gamma_t : A \rightarrow B$, $t \in [0, 1]$ such that $t \mapsto \gamma_t(a)$ is norm continuous for all $a \in A$ and with $\gamma_0 = \alpha$ and $\gamma_1 = \beta$), then $\alpha_* = \beta_*$.
- **Stable** For any C^* -algebra A and any rank 1 projection $p \in \mathcal{K}(\mathcal{H})$, the morphism $a \mapsto a \otimes p$ from A to $A \otimes \mathcal{K}(\mathcal{H})$ induces an isomorphism from $F(A)$ onto $F(A \otimes \mathcal{K}(\mathcal{H}))$.
- **Continuous** if whenever $\{A_n, \phi_n\}_{n=1}^\infty$ is a countable directed sequence, then $F(\varinjlim(A_n, \phi_n)) = \varinjlim(F(A_n), \phi_{n*})$

K_0 and K_1 are half exact, homotopy invariant, stable, and continuous.

Theorem: If F is a functor that is half exact, homotopy invariant, stable, and continuous with $F(\mathbb{C}) = \mathbb{Z}$ and $F(S\mathbb{C}) = 0$ then F is K_0 .

Theorem: If F is a functor that is half exact, homotopy invariant, stable, and continuous with $F(\mathbb{C}) = 0$ and $F(S\mathbb{C}) = \mathbb{Z}$ then F is K_1 .

Other K -theory Results

Other K -theory Results

Direct Sums: If A and B are C^* -algebras, then

$$K_0(A \oplus B) \cong K_0(A) \oplus K_0(B) \quad \text{and} \quad K_1(A \oplus B) \cong K_1(A) \oplus K_1(B).$$

Other K -theory Results

Direct Sums: If A and B are C^* -algebras, then

$$K_0(A \oplus B) \cong K_0(A) \oplus K_0(B) \quad \text{and} \quad K_1(A \oplus B) \cong K_1(A) \oplus K_1(B).$$

Split exact sequences: If we have a split exact sequence

$$0 \longrightarrow I \xrightarrow{i} A \xrightleftharpoons[\pi]{s} A/I \longrightarrow 0$$

then K_0 and K_1 each take it to a split exact sequence

$$0 \longrightarrow K_0(I) \xrightarrow{i_0} K_0(A) \xrightleftharpoons[\pi_0]{s_0} K_0(A/I) \longrightarrow 0 \quad 0 \longrightarrow K_1(I) \xrightarrow{i_1} K_1(A) \xrightleftharpoons[\pi_1]{s_1} K_1(A/I) \longrightarrow 0$$

Other K -theory Results

Direct Sums: If A and B are C^* -algebras, then

$$K_0(A \oplus B) \cong K_0(A) \oplus K_0(B) \quad \text{and} \quad K_1(A \oplus B) \cong K_1(A) \oplus K_1(B).$$

Split exact sequences: If we have a split exact sequence

$$0 \longrightarrow I \xrightarrow{i} A \xleftarrow[\pi]{s} A/I \longrightarrow 0$$

then K_0 and K_1 each take it to a split exact sequence

$$0 \longrightarrow K_0(I) \xrightarrow{i_0} K_0(A) \xleftarrow[\pi_0]{s_0} K_0(A/I) \longrightarrow 0 \quad 0 \longrightarrow K_1(I) \xrightarrow{i_1} K_1(A) \xleftarrow[\pi_1]{s_1} K_1(A/I) \longrightarrow 0$$

Tensor Products: The Künneth Theorem says that if A and B are nuclear and their K -groups are all torsion free, then

$$K_0(A \otimes B) \cong (K_0(A) \otimes K_0(B)) \oplus (K_1(A) \otimes K_1(B))$$

$$K_1(A \otimes B) \cong (K_0(A) \otimes K_1(B)) \oplus (K_1(A) \otimes K_0(B))$$

Pimsner-Voiculescu Exact Sequence for crossed products by \mathbb{Z}

If A is a unital C^* -algebra and α is a $*$ -automorphism of A , we may form the crossed product $A \times_{\alpha} \mathbb{Z}$. If we let $i : A \hookrightarrow A \times_{\alpha} \mathbb{Z}$ denote the natural embedding, then there is an exact sequence

$$\begin{array}{ccccc} K_0(A) & \xrightarrow{id-\alpha_0} & K_0(A) & \xrightarrow{i_0} & K_0(A \times_{\alpha} \mathbb{Z}) \\ \uparrow & & & & \downarrow \\ K_1(A \times_{\alpha} \mathbb{Z}) & \xleftarrow{i_1} & K_1(A) & \xleftarrow{id-\alpha_1} & K_1(A) \end{array}$$

Note: This 6-term sequence does *not* come from a short exact sequence.

Pimsner-Voiculescu Exact Sequence for crossed products by \mathbb{Z}

If A is a unital C^* -algebra and α is a $*$ -automorphism of A , we may form the crossed product $A \times_{\alpha} \mathbb{Z}$. If we let $i : A \hookrightarrow A \times_{\alpha} \mathbb{Z}$ denote the natural embedding, then there is an exact sequence

$$\begin{array}{ccccc}
 K_0(A) & \xrightarrow{id-\alpha_0} & K_0(A) & \xrightarrow{i_0} & K_0(A \times_{\alpha} \mathbb{Z}) \\
 \uparrow & & & & \downarrow \\
 K_1(A \times_{\alpha} \mathbb{Z}) & \xleftarrow{i_1} & K_1(A) & \xleftarrow{id-\alpha_1} & K_1(A)
 \end{array}$$

Note: This 6-term sequence does *not* come from a short exact sequence.

Application: If A is an $n \times n$ matrix and \mathcal{O}_A is the associated Cuntz-Krieger algebra, (a dual version of) the above sequence can be used to obtain

$$\begin{array}{ccccc}
 \mathbb{Z}^n & \xrightarrow{I-A^t} & \mathbb{Z}^n & \longrightarrow & K_0(\mathcal{O}_A) \\
 \uparrow & & & & \downarrow \\
 K_1(\mathcal{O}_A) & \longleftarrow & 0 & \longleftarrow & 0
 \end{array}$$

So $K_0(\mathcal{O}_A) \cong \text{coker}(I - A^t)$ and $K_1(\mathcal{O}_A) \cong \ker(I - A^t)$.

Relation with Topological K -theory

If X is a compact Hausdorff space, the n^{th} topological K -group of X is isomorphic to $K_n(C(X))$.

AF-algebras

If A is an AF-algebra, $A = \varinjlim(A_n, \phi_n)$, with each A_n finite-dimensional. Thus each A_n is a direct sum of matrix algebras, and by the continuity of K -theory and the fact K -theory distributes over direct sums

$$K_0(A) = \varinjlim(K_0(A_n), (i_n)_0) = \varinjlim(K_0(A_n), (i_n)_0) = \varinjlim(\mathbb{Z}^{k_n}, (i_n)_0)$$

and

$$K_1(A) = \varinjlim(K_1(A_n), (i_n)_1) = \varinjlim(0, (i_n)_1) = \{0\}.$$

Therefore, when A is an AF-algebra, $K_1(A) = 0$. Also, $K_0(A)$ is a direct limit of \mathbb{Z}^{n_k} 's and, in particular, $K_0(A)$ has no torsion.

Stabilization and Morita Equivalence

A C^* -algebra is **stable** if $A \otimes \mathcal{K}(\mathcal{H}) \cong A$.

Stabilization and Morita Equivalence

A C^* -algebra is **stable** if $A \otimes \mathcal{K}(\mathcal{H}) \cong A$.

For any C^* -algebra A , the **stabilization of A** is defined to be $A \otimes \mathcal{K}(\mathcal{H})$. The stabilization $A \otimes \mathcal{K}(\mathcal{H})$ is stable because $\mathcal{K}(\mathcal{H}) \otimes \mathcal{K}(\mathcal{H}) \cong \mathcal{K}(\mathcal{H})$, so

$$(A \otimes \mathcal{K}(\mathcal{H})) \otimes \mathcal{K}(\mathcal{H}) \cong A \otimes (\mathcal{K}(\mathcal{H}) \otimes \mathcal{K}(\mathcal{H})) \cong A \otimes \mathcal{K}(\mathcal{H}).$$

Another way to view the stabilization: Since $\overline{M_\infty(\mathbb{C})} = \mathcal{K}(\mathcal{H})$, we have

$$A \otimes \mathcal{K}(\mathcal{H}) \cong A \otimes \overline{M_\infty(\mathbb{C})} \cong \overline{A \otimes M_\infty(\mathbb{C})} \cong \overline{M_\infty(A)}.$$

We say A and B are **stably isomorphic** when $A \otimes \mathcal{K}(\mathcal{H}) \cong B \otimes \mathcal{K}(\mathcal{H})$

Theorem: If A and B have countable approximate units (e.g., they are unital or separable), then A and B are Morita equivalent if and only if A and B are stably isomorphic.

K -theory as an Invariant

Our groups K_0 and K_1 are stable:

$$K_0(A) \cong K_0(M_n(A)) \cong K_0(A \otimes \mathcal{K}(\mathcal{H}))$$

$$K_1(A) \cong K_1(M_n(A)) \cong K_1(A \otimes \mathcal{K}(\mathcal{H}))$$

Thus K -theory only “sees” a C^* -algebra up to Morita equivalence; i.e., if A and B are Morita equivalent, then $K_0(A) \cong K_0(B)$ and $K_1(A) \cong K_1(B)$. In other words, K -theory is a Morita equivalence invariant.

K -theory as an Invariant

Our groups K_0 and K_1 are stable:

$$K_0(A) \cong K_0(M_n(A)) \cong K_0(A \otimes \mathcal{K}(\mathcal{H}))$$

$$K_1(A) \cong K_1(M_n(A)) \cong K_1(A \otimes \mathcal{K}(\mathcal{H}))$$

Thus K -theory only “sees” a C^* -algebra up to Morita equivalence; i.e., if A and B are Morita equivalent, then $K_0(A) \cong K_0(B)$ and $K_1(A) \cong K_1(B)$. In other words, K -theory is a Morita equivalence invariant.

K -theory can therefore be used to show two C^* -algebras are “different”, where “different” means “not Morita equivalent”. For example,

$$K_0(\mathcal{O}_n) \cong \mathbb{Z}/n\mathbb{Z}.$$

Thus the Cuntz algebra \mathcal{O}_n is not Morita equivalent to \mathcal{O}_m when $n \neq m$.

K -theory as an Invariant

Our groups K_0 and K_1 are stable:

$$K_0(A) \cong K_0(M_n(A)) \cong K_0(A \otimes \mathcal{K}(\mathcal{H}))$$

$$K_1(A) \cong K_1(M_n(A)) \cong K_1(A \otimes \mathcal{K}(\mathcal{H}))$$

Thus K -theory only “sees” a C^* -algebra up to Morita equivalence; i.e., if A and B are Morita equivalent, then $K_0(A) \cong K_0(B)$ and $K_1(A) \cong K_1(B)$. In other words, K -theory is a Morita equivalence invariant.

K -theory can therefore be used to show two C^* -algebras are “different”, where “different” means “not Morita equivalent”. For example,

$$K_0(\mathcal{O}_n) \cong \mathbb{Z}/n\mathbb{Z}.$$

Thus the Cuntz algebra \mathcal{O}_n is not Morita equivalent to \mathcal{O}_m when $n \neq m$.

In some cases, K -theory can also be used to show two C^* -algebras are “the same”, where “the same” sometimes means “Morita equivalent” and sometimes means “isomorphic”. In these situations, we say K -theory is a complete invariant.

Classification of AF-algebras

Let A be an AF-algebra. Recall $K_1(A) = 0$, so all K -theory info is in the K_0 -group. Since A has a countable approximate unit of projections,

$$K_0(A) = \{[p] - [q] : p, q \in \text{Proj } M_\infty(A)\}.$$

We define the **positive elements** of $K_0(A)$ to be

$$K_0(A)^+ = \{[p] : p \in \text{Proj } M_\infty(A)\}.$$

Defining $a \leq b$ iff $b - a \in K_0(A)^+$ gives a partial ordering on $K_0(A)$.

We define the **scale** of $K_0(A)$ to be

$$\Sigma(A) = \{[p] : p \in \text{Proj}(A)\}.$$

Theorem (Elliott)

Let A and B be AF-algebras.

(1) A is Morita equivalent to B iff $(K_0(A), K_0(A)^+) \cong (K_0(B), K_0(B)^+)$.

(2) $A \cong B$ iff $(K_0(A), K_0(A)^+, \Sigma(A)) \cong (K_0(B), K_0(B)^+, \Sigma(B))$.

Moreover, when A (respectively, B) is unital, we may replace $\Sigma(A)$ by $[1_A]$ (respectively, we may replace $\Sigma(B)$ by $[1_B]$).

Classification of Purely Infinite, Simple C^* -algebras

Let A be a C^* -algebra that is purely infinite and simple. Then

$K_0(A) = K_0(A)^+ = \{[p] : p \in \text{Proj } M_\infty(A)\}$. If A is also unital, then

$K_0(A) = \Sigma(A) = \{[p] : p \in \text{Proj}(A)\}$.

Classification of Purely Infinite, Simple C^* -algebras

Let A be a C^* -algebra that is purely infinite and simple. Then $K_0(A) = K_0(A)^+ = \{[p] : p \in \text{Proj } M_\infty(A)\}$. If A is also unital, then $K_0(A) = \Sigma(A) = \{[p] : p \in \text{Proj}(A)\}$.

Theorem (Kirchberg and Phillips)

Let A and B be purely infinite, simple C^* -algebras that are also separable and nuclear.¹

(1) If A and B are nonunital, the following are equivalent:

- (a) A is Morita equivalent to B .
- (b) A is isomorphic to B .
- (c) $K_0(A) \cong K_0(B)$ and $K_1(A) \cong K_1(B)$.

(2) If A and B are unital, then

- (i) A is Morita equivalent to B iff $K_0(A) \cong K_0(B)$ and $K_1(A) \cong K_1(B)$.
- (ii) A is isomorphic to B iff $(K_0(A), [1_A]) \cong (K_0(B), [1_B])$ and $K_1(A) \cong K_1(B)$.

¹Technically, we also need A and B to be in the bootstrap class to which the UCT applies, but let's not get into that.

Classification of simple nuclear C^* -algebras

Elliott conjectured that all simple, separable, nuclear C^* -algebras can be classified up to Morita equivalence by an invariant $\text{Ell}(A)$ that includes the ordered K_0 -group, the K_1 -group, and other data provided by K -theory.

¹To be more precise: (1) \iff (2) has been established and (1) \iff (2) \iff (3) is known in many cases (e.g., when the trace space of the C^* -algebra has finitely many extreme points) but has yet to be proven in general.

Classification of simple nuclear C^* -algebras

Elliott conjectured that all simple, separable, nuclear C^* -algebras can be classified up to Morita equivalence by an invariant $\text{Ell}(A)$ that includes the ordered K_0 -group, the K_1 -group, and other data provided by K -theory.

Counterexamples showed the conjecture is not true for *all* simple, separable, nuclear C^* -algebras — one needs an additional hypothesis, which may be formulated in various ways. TFAE:

- (i) A has finite nuclear dimension.
- (ii) A is \mathcal{Z} -stable; i.e., $A \cong A \otimes \mathcal{Z}$ where \mathcal{Z} is the Jiang-Su algebra.
- (iii) A has strict comparison of positive elements.¹

¹To be more precise: (1) \iff (2) has been established and (1) \iff (2) \iff (3) is known in many cases (e.g., when the trace space of the C^* -algebra has finitely many extreme points) but has yet to be proven in general.

Classification of simple nuclear C^* -algebras

Elliott conjectured that all simple, separable, nuclear C^* -algebras can be classified up to Morita equivalence by an invariant $\text{Ell}(A)$ that includes the ordered K_0 -group, the K_1 -group, and other data provided by K -theory.

Counterexamples showed the conjecture is not true for *all* simple, separable, nuclear C^* -algebras — one needs an additional hypothesis, which may be formulated in various ways. TFAE:

- (i) A has finite nuclear dimension.
- (ii) A is \mathcal{Z} -stable; i.e., $A \cong A \otimes \mathcal{Z}$ where \mathcal{Z} is the Jiang-Su algebra.
- (iii) A has strict comparison of positive elements.¹

Theorem (By many hands)

Let A and B be simple, separable, nuclear C^ -algebras satisfying one (and hence all) of the above three conditions. Then $A \cong B$ if and only if $\text{Ell}(A) \cong \text{Ell}(B)$.*

¹To be more precise: (1) \iff (2) has been established and (1) \iff (2) \iff (3) is known in many cases (e.g., when the trace space of the C^* -algebra has finitely many extreme points) but has yet to be proven in general.

What about non-simple C^* -algebras?

What about non-simple C^* -algebras?

Elliott's Theorem applies to non-simple AF-algebras. Some progress has also been made for purely infinite C^* -algebras.

What about non-simple C^* -algebras?

Elliott's Theorem applies to non-simple AF-algebras. Some progress has also been made for purely infinite C^* -algebras.

Far-reaching results have also been obtained for graph C^* -algebras (which contain the Cuntz-Krieger algebras and the AF-algebras as subclasses).

What about non-simple C^* -algebras?

Elliott's Theorem applies to non-simple AF-algebras. Some progress has also been made for purely infinite C^* -algebras.

Far-reaching results have also been obtained for graph C^* -algebras (which contain the Cuntz-Krieger algebras and the AF-algebras as subclasses).

Theorem (Eilers and T)

Let A be a separable graph C^ -algebra with exactly one ideal I . Then A is classified up to Morita equivalence by the 6-term exact sequence*

$$\begin{array}{ccccc} K_0(I) & \xrightarrow{i_0} & K_0(A) & \xrightarrow{\pi_0} & K_0(A/I) \\ \delta_1 \uparrow & & & & \downarrow \delta_0 \\ K_1(A/I) & \xleftarrow{\pi_1} & K_1(A) & \xleftarrow{i_1} & K_1(I) \end{array}$$

where the K_0 -groups in the invariant are considered as ordered groups.

A complete classification up to Morita equivalence has been obtained for C^* -algebras of finite graphs.

The invariant, called [ordered, filtered \$K\$ -theory](#) includes the 6-term exact sequences of every ideal and subquotient of A .

A complete classification up to Morita equivalence has been obtained for C^* -algebras of finite graphs.

The invariant, called [ordered, filtered \$K\$ -theory](#) includes the 6-term exact sequences of every ideal and subquotient of A .

Theorem (Eilers, Restorff, Ruiz, and Sorensen)

Let A be a separable graph C^ -algebra of a finite graph. Then A is classified up to Morita equivalence by its ordered, filtered K -theory.*

Generalizations of K -theory

Using extensions, it is possible to create a contravariant theory, called K -homology that assigns groups $K^0(A)$ and $K^1(A)$ to a C^* -algebra A .

KK -theory is a bivariate functor that takes a pair of C^* -algebra (A, B) and assigns an abelian group $KK(A, B)$.

It turns out that

- $KK(\mathbb{C}, A) \cong K_0(A)$
- $KK(S\mathbb{C}, A) \cong K_1(A)$
- $KK(A, \mathbb{C}) \cong K^0(A)$
- $KK(A, S\mathbb{C}) \cong K^1(A)$

Recall: $S\mathbb{C} = C_0(\mathbb{R})$.

So KK -theory simultaneously generalizes K -theory and K -homology, and can be viewed as a bivariate pairing between the two theories.

There is also a variant of KK -theory, known as E -theory, that was developed to get more (and better) exact sequences.

Table of K -groups

A	$K_0(A)$	$K_1(A)$
\mathbb{C}	\mathbb{Z}	0
M_n	\mathbb{Z}	0
\mathbb{K}	\mathbb{Z}	0
\mathbb{B}	0	0
\mathbb{B}/\mathbb{K}	0	\mathbb{Z}
$C_0(\mathbb{R}^{2n})$	\mathbb{Z}	0
$C_0(\mathbb{R}^{2n+1})$	0	\mathbb{Z}
$C(\mathbb{T}^n)$	$\mathbb{Z}^{2^{n-1}}$	$\mathbb{Z}^{2^{n-1}}$
$C(S^{2n})$	\mathbb{Z}^2	0
$C(S^{2n+1})$	\mathbb{Z}	\mathbb{Z}
\mathcal{T}	\mathbb{Z}	0
\mathcal{O}_n	$\mathbb{Z}/(n-1)$	0
A_θ	\mathbb{Z}^2	\mathbb{Z}^2
II_1 -factor	\mathbb{R}	0

To learn more about K -theory, visit your local library . . .

Introductory Textbooks

- “ K -theory and C^* -algebras. A friendly approach” by N.E. Wegge-Olsen.
- “An introduction to K -theory for C^* -algebras” by M. Rørdam, F. Larsen, and N. Laustsen

Harder Textbook

- “ K -theory for operator algebras”, Second Edition, by B. Blackadar

A crash course on the K_0 -group and Elliott’s theorem for AF-algebras appears in Sec. III and Sec. IV of Davidson’s book.

- “ C^* -algebras by example” by K. Davidson.

